

## LOPRO OR GVL STAT


1 Insert the blade midline, advancing the tip until it is past the posterior portion of the tongue.




2 Identify the epiglottis and advance the blade into the vallecula for a view of the larynx.




3 Introduce the ET tube along the right side the blade.



4 Complete the intubation.



5 To remove the GlideRite stylet, pull it out towards the patient's feet.





– Look in the mouth during step.



Look at the monitor during step.





\tag{Important: This document is for quick reference only. For complete instructions, see your Operations & Maintenance Manual at verathon.com/product-documentation.

Verathon Medical (Canada) ULC 2227 Douglas Road Burnaby, BC V5C 5A9 Canada

Tel: +1 604 439 3009 Fax: +1 604 439 3039 EC REP

Verathon Medical (Europe) B.V. Willem Fenengastraat 13 1096 Bl. Amsterdam, The Netherlands Tel: +31 (0) 20 210 30 91 Fax: +31 (0) 20 210 30 92 Verathon Medical (Australia) Pty Limited Unit 9, 39 Herbert Street St Leonards NSW 2065

Australia Within Australia: 1800 613 603 Tel / 180

1800 613 603 Tel / 1800 657 970 Fax International:

+61 2 9431 2000 Tel / +61 2 9475 1201 Fax



